

Photosynthesis Components *in vitro* and *ex vitro* Date Palm

Darwesh, Rasmia, S.S.Central Lab. For Research and Development
of Date Palm (ARC) Giza, Egypt**Hemmat, A. Ibrahim**Agric. Biochem. Dept., Fac of Agric.,
Ain Shams Univ., Egypt**Hany, A. Mahmoud**Agric. Biochem. Dept., Fac of Agric.,
Ain Shams Univ., Egypt

Email ID : darweshsrasmia@gmail.com

Abstract – Date palm (*Phoenix dactylifera* L.) trees one of the important popular fruits were found in all Egypt cultivation areas, *in vitro* technique is a control condition as temperature, water, light and carbohydrates which can affect plants photosynthesis processes and own production as, pigments, proteins, indoles and enzymes, *ex vitro* plantlets can be modify these effects, thus this work investigate effects of *in vitro* and *ex vitro* date palm on these components, the analysis of tissues *in vitro* shooting and rooting stages showed smallest contents of pigments, proteins and indoles, the antioxidant enzymes CAT and POD activities increased progressively from *in vitro* shooting and rooting stages to *ex vitro* plantlets in the greenhouse and open field in addition to offshoot correlated to the growth stages and changes in the environmental condition. In the C4 plants photosynthetic activities are take place between mesophyll and bundle sheath cells where the initial carboxylating enzyme is PEPC which greater in the offshoots and plantlets in the open field than *in vitro* tissues that have the lowest activity enzyme as well as Rubisco enzyme in the tested tissues assumed gradual increasing from *in vitro* to *ex vitro* in the greenhouse and open field.

Keywords – Photosynthesis Activity, Date Palm, Enzymes, Indoles, Pigments, Proteins.

I. INTRODUCTION

Date palm (*Phoenix dactylifera* L.) is a monocotyledonous, angiosperm and dioecious trees have specific carbon – metabolism to produce different organic contents. Date palm tissue culture technique has provide large numbers of fruiting trees under specific condition as constant light, temperature, water also sufficient minerals and carbohydrates, this *in vitro* control condition continuous in all stages of tissue culture until the plantlets transferred to acclimatization stage in the greenhouse which facing different condition caused failure or decreasing success of *ex vitro* acclimatization, [1] Darwesh 2015, therefore, many scientists investigate the physiology characters *in vitro* and *ex vitro* as photosynthesis, enzymes, pigments and proteins. High air humidity, low gas exchange and thus a CO₂-shortage in the whole photoperiod, and relatively low photosynthetic photon flux density (PPFD), induce disturbances in plant development and photosynthetic performance [2] Pospisilova *et al.* 1997 and [3] Kozai *et al.* 1997, Rubisco content was substantially decreased in *in vitro* leaves under elevated CO₂ (1000 μmol.mol⁻¹) was supplied [4] De La Vina *et al.* 1999, plantlets exhibit abnormal morphology, anatomy, and physiology as a response of *in vitro* condition as high air humidity, low irradiation, low CO₂ and high concentration of sugars as carbon source [5] Desjardins 1995 and [6]

Kozai and Smith 1995, *In vitro* avocado, oak, and strawberry showed a decrease of Rubisco and significant decrease in total soluble proteins of avocado plantlets [7] Rival *et al.* 1999 on coconut and oil palm and [8] Premkumar *et al.* 2001, Moreover, these bad effects of *in vitro* can be changes in *ex vitro* greenhouse, that causing increasing of photosynthesis components as enzymes activity, proteins and chlorophyll contents, Increasing CO₂ level produced increasing photosynthesis rates also chlorophyll, carbohydrates contents and Rubisco activity [9] Cournac *et al.* 1992 and [10] Kadlecck *et al.* 2001, *Spathiphyllum floribundum* plantlets grown under *in vitro* conditions have differences in net photosynthetic rate, chl, and Fv/Fm which disappeared during the 15 days of *ex vitro* acclimatization and increase in net photosynthetic rate [11] Van Huylenbroeck, and Debergh 1996, *in vitro* of avocado (*Persea americana* Mill.) with highest concentration of sucrose showed significant decreasing of maximum photosynthetic rate, carbohydrates and starch [12] Ticha *et al.* 1998, significant rise in chlorophylls in *ex vitro* grapevine leaves after transplantation [13] Amancio *et al.* 1999, C3 plants found in cool and wet climates, with low light intensity, while C4 plants occur in hot and dry climatic conditions with high light intensity, C4 and CAM plants usually in arid environments. C4 plants have higher photosynthetic efficiency than C3 plants in arid, hot, and under high-light conditions, they possess an additional carbon fixation pathway and limit photorespiration [14] Taiz and Zeiger 2010, in C4 plant phosphoenolpyruvate carboxylase (PEPC) is the enzyme responsible for catalyzing the primary fixation of atmospheric CO₂, PEPC kinase (PEPCK), a protein kinase involved in this phosphorylation of PEPC [15] Agetsuma *et al.* 2005 which converted to oxaloacetate through phosphoenolpyruvate and carbon dioxide reaction [16] Méndez-Lucas *et al.* 2014, oxaloacetate converted to aspartate, which move to bundle sheath, in the bundle sheath cells, aspartate is converted back to oxaloacetate. PEPCK decarboxylates the bundle sheath oxaloacetate, releasing carbon dioxide, which fixed by the enzyme Rubisco [17] Christopher and Holtum 1996 and [18] Kanai and Edwards 1999, photosynthesis rate and growth were decreased of *in vitro* plantlets (*Samanea saman* Merr) and their enhancing after transfer to the greenhouse [19] Kriengkrai *et al.* 2004, the main bundle fibrous sheath thick of date palm leaves was different between different soft and dry cultivars [20] Abd El-Baky 2012, an incessant activity of Catalase, Ascorbate Peroxidase and Glutathione Reductase were also detected through one week of acclimatization of *Cardiospermum halicacabum* plantlets [21] Anushi and Anis 2014. The

present work was done in order to investigate the effect of *in vitro* conditions on the photosynthesis components (pigments, proteins, indoles and enzymes) and changes after the plantlets transferring to *ex vitro* under different conditions in the greenhouse.

II. MATERIALS AND METHODS

This work was done in 2015- 2016 in the Central Laboratory for Research and Development of date palm and Biochem. Dept., Fac of Agric., Ain Shams Univ. to measure photosynthesis components of date palm (*Phoenix dactylifera* L.) *in vitro* and *ex vitro*. *In vitro* leaves date palm samples were taken from shooting and rooting stages (cultured in MS media supplemented with 0.1 mg/l NAA + 0.05 BA + 30 g/l sucrose) placed in a growth chamber (27± C⁰, 7000 lux), samples were taken as

1. *in vitro* shooting stage
2. *in vitro* rooting stage
3. *ex vitro* plantlets greenhouse (6 months from acclimatization stage)
4. *ex vitro* plantlets open field (one year from acclimatization stage)
5. offshoots

Samples *in vitro* and *ex vitro* were prepared to measure the following estimations

1. **Chlorophyll contents** a, b, total chlorophyll and carotenoids mg/g f.wt.: as described by [22] **Lichtenthaler and Wellburn** 1994
2. **Proteins** mg/g f.wt.: as described by [23] **Bradford** 1976
3. **Indoles** mg/g f.wt.: as [24] **Larsen et al.** 1962
Antioxidant Enzyme Activity CAT and POD
Catalase CAT (EC 1.11.1.6) in leaves µmol /g.f.wt./min: as [25] **Aebi**, 1984
Peroxidase POD (PRX, E.C.1.11.1.7) in leaves µmol /g.f.wt./min: as [26] **Polle et al.**, 1994
4. **Phosphoenolpyruvate carboxykinase** namely **PEPC** (EC 4.1.1.31) Unit /mg protein: as [27] **Gonzalez** 1998
5. **Ribulose1, 5-bisphosphate carboxylase/oxygenase** (**Rubisco**) (EC 4.1.1.39) Unit /mg protein: as [28] **Lilley** 1974

Experimental design: Complete randomized block design with three replicates and three plantlets for each one, two growth seasons (8 months for each). Data were analyzed by analysis of variances (ANOVA) and the means were compared following t- test using L.S.D. values at 5 % level [29] **Snedecor and Chocran** 1990).

III. RESULTS

3.1. Leaves chlorophyll a, b and total chlorophyll mg/g f.w.: determined of chlorophyll a, b as well as total chlorophyll (Fig 1 and 7) on the different types tissues *in vitro* and *ex vitro* produce the evidence that the chlorophyll a, b and total chlorophyll were increased significantly from shooting tissues in the shooting stage as 0.13 mg/g f.wt. for chl. a, 0.15 mg/g f.wt. for rooting tissues, 0.2 mg/g f.wt. for *ex vitro* plantlets in

the greenhouse and have the insignificant differs 0.4 mg/g f.wt. between *ex vitro* plantlets in the open field and offshoots. Assayed chlorophyll b showed significant gradually raising from shooting tissues 0.02 mg/g f.wt., 0.13 mg/g f.wt. for rooting tissues, 0.12 mg/g f.w. for *ex vitro* plantlets greenhouse, 0.2 mg/g f.wt. for *ex vitro* plantlets in the open field and offshoots with insignificant variance in between. Consequent results found on the total chlorophyll produced from *in vitro* shooting tissues 0.15mg/g f.wt., 0.25 mg/g f.wt. for *in vitro* rooting stage, 0.32 mg/g f.wt. for *ex vitro* greenhouse, insignificant distinction 0.6 mg/g f.wt. obtained between *ex vitro* plantlets open field and offshoots

3.2. Proteins contents mg/g f.wt.: different types of tissues *in vitro* and *ex vitro* (Fig 2 and 7) presented significantly higher different values, smaller values of proteins associated to *in vitro* shooting tissues 1.1 mg/g f.wt. which ascending by rooting tissues 2.2 mg/g f.wt., moreover *ex vitro* tissues exhibited significant increasing of proteins 2.3 mg/g f.wt. for plantlets greenhouse, the plantlets in the open field and offshoots produced insignificant distinguish in between 2.8 and 2.9 mg/g f.wt. respectively

3.3. Indoles contents mg/g f.wt.: *in vitro* shooting tissues (Fig 3 and 7) recorded smallest indole contents 3.6 mg/g f.wt. followed by *in vitro* tissue of rooting 10.2 mg/g f.wt. meanwhile *ex vitro* plantlet in the open field and offshoots recorded highest indole contents 11.5 and 12.1 mg/g f.wt. respectively without significant differs between them

3.4. Antioxidant enzymes CAT and POD µmol/ g. f.wt./min: increasing activity of antioxidant enzymes CAT and POD (Fig 4 and 7) significantly pronounced enhancing at the *ex vitro* tissues in the greenhouse and open field meanwhile the assaying enzymes activity in the *in vitro* tissues shooting and rooting stages gave minimum activity. non- significant differences were found for enzymes activity between plantlets in the open field and offshoot for two enzymes.

3.5. PEPC and Rubisco enzymes : spectrophotometer data on the determination of photosynthesis enzymes (Figs 5, 6 and 7) Phosphoenolpyruvate carboxykinase namely PEPC (EC 4.1.1.31) and ribulose 1, 5-biphosphate carboxylase-oxygenase (EC 4.1.1.39; Rubisco) showed significant variance from *in vitro* and *ex vitro* tissues, the significant differences were observed between the different types *in vitro* and *ex vitro* tissues related to PEPC and Rubisco activity which recorded highest activity for offshoots meanwhile *in vitro* tissues bringing out lowest activity followed by rooting stage, greenhouse and open field.

IV. DISCUSSION

Date palm (*Phoenix dactylifera* L.) is the important genus in Arecaceae distributed in the arid and semi arid area [30] Jain 2012 and the main source of food in these regions of North Africa, middle east and South-Asian countries [31]

FAO 2013, date palm is dioecious species and very heterozygous separate male and female palms, propagation by seedlings produced over 50% male (one male tree for pollen 25 female trees) in addition male and female seedlings can be distinction flowering stage, offshoots as a traditional propagation of date palm produced limited numbers of trees which insufficient for new reclamation areas, the survival rate of offshoots is low when separated from mother tree by incorrectly practice or infection by palm diseases and pests, thus in vitro micropropagation become the most suitable for providing great numbers of homogenous and healthy female date palm in this respect many researchers as [32] Eke et al. 2005, [33] Al-Mazroui et al., 2007, [34] Al- Khatieb 2008b, [35] Mustafa et al. 2013 and [36] Mazri and Meziani 2015, on the other hand in vitro culture control conditions resulted in abnormal morphology, anatomy and physiology of plantlets and might be impaired many photosynthesis functions thus needed to acclimatization stage to correct these abnormalities, after transfer from the in vitro to the ex vitro greenhouse [37] Fila et al. 1998, [38] Bolar et al. 1998. **In concerning to leaves contents of chlorophyll, proteins and indoles**, chlorophyll can play a vital role for building growth components as proteins, auxins etc., results showed the progressively increasing of chlorophyll a and b also total chlorophyll as well as increasing of proteins and indoles contents, increase in chlorophyll a and b from in vitro to ex vitro of date palm plantlets demonstrated the induction of chlorophyll synthesis enzymes indispensable for chlorophyll biosynthesis, increase in Chlorophyll a and b synchronous with the new leaves during the acclimatization, Chlorophyll *a* fluorescence has been used as an indicator of photochemical efficiency [39] Krause and Weis, 1999, Leaves formed during the acclimatization period may still have a lower photosynthetic capacity than leaves of greenhouse-grown plants [40] Carvalho et al. 2001, decreasing chlorophyll of Rhododendron 'Alfred' value and increasing vitality index value (Rfd) during first 4 weeks of acclimatization plantlets proved changing from mixotrophic metabolism of plantlets *in vitro* to autotrophic in ex vitro [41] Bošena Matysiak 2004, growing plants maintain the optimum nutrient levels that led to high chlorophyll activity which may be correlate with high amount of protein contents [42] Arigita et al. 2005, higher photosynthetic rates P(N) of sun leaves (ginkgo and beech) and stomatal conductance were also reflected by higher values for the Chl fluorescence decrease ratios. [43] Lichtenthaler et al. 2007 and [44] 2013, chlorophyll a, b and total chlorophyll increased in hardening and acclimatization stages of *Boswellia serrate* Roxb. Plantlets [45] Suthar and Purohit 2011, ex vitro plants have greater chlorophyll a and b contents of *Nepeta nuda* ssp. *nuda* than in vitro plants [46] Dragolova et al. 2015, chlorophyll and total soluble proteins were increased in the glasshouse in the acclimatization stage of *Tectona grandis* L. [47] Akram and Aftab 2015, increasing protein and chlorophyll contents at the end of acclimatization period [48] Hofman et al. 2002, protein content increased at the end of shoot formation stage [49] Sharifi 2012, indoles contents

increased at the end of pre-acclimatization stage [50] Darwesh et al. 2011, and increasing of chlorophyll a and b and indoles contents in acclimatization stage was found in date palm plantlets in the greenhouse [51] Ibrahim et al. 2012.

Activity of antioxidant enzymes: assaying CAT and POD activity showed increasing activity sequenced from in vitro shooting stage to ex vitro plantlets in the greenhouse and in the open field and offshoots which come out from the differs of environmental conditions from in vitro and ex vitro also different growth stages with different photosynthesis products, antioxidants helped in preventing the damage occurred in cells by free radicals that are released during normal metabolic process of oxidation [52] Young and Woodside 2001, antioxidant activity of polyphenols due to redox properties which are important in decomposing peroxides [53] Karou et al., 2005, superoxide dismutase and guaiacol peroxidase activities increased with increasing light intensity, Catalase and glutathione reductase increased during the first three weeks of acclimatization, while an increase in guaiacol peroxidase and ascorbate peroxidase observed later in the acclimatization, [54] Van Huylbroeck et al. 2000, rise activity of SOD after one week of transplantation plantlets showed its role in preventing oxidative stress of *Rauvolfia tetraphylla*, increasing carotenoids content reflects the functional response of photosynthetic apparatus to the different light environment of *Tylophora indica* [55] Faisal and Anis 2009 and [56] 2015, catalase and glutathione peroxidase plays an important role in preventing cell injuries [57] Starlin and Gopalakrishnan, 2013, CAT decomposition of H₂O₂ into oxygen and water associated with guard systems for AOS in cells [58] Agnieszka and Edyta 2014, the activity of catalase and peroxidase changes with fluctuations of endogenous H₂O₂ content [59] Konieczny et al. 2014.

PEPC and Rubisco enzymes: findings on PEPC and Rubisco activity showed graduated value activity from date palm in vitro tissues which scored lowest activity values to maximum activity values belonging to ex vitro tissues. C3 and C4 plants use photosynthesis process to convert light to energy and CO₂ into (carbohydrates). increase of both biomass and CO₂ fixation with light intensity in C4 faster than C3, led to use of solar energy in C4 plants, C4 plants following angiosperms and monocotyledon with high daytime temperatures, intense sunlight, C4 plants can grow well at 35-40 °C, grow in 20-25 °C soil temperature [60] Oberhuber et al. 1993 and [61] Wang et al. 2012, Date palm can assimilate CO₂ through two independent carboxylation pathways, one occurs in the chloroplasts through, another pathway occurs in the cytosol through the Phosphoenolpyruvate carboxylase pathway [62] Kwa et al., 1997, C4 plants, CO₂ is fixed by phosphoenolpyruvate carboxylase (PEPC) in the mesophyll cells into four-carbon acids and flow into an inner ring of bundle sheath cells, which decarboxylated and the CO₂ is fixed again by Rubisco, [63] Doubnerová and Ryšlavá 2011, [64] Freschi and Mercier 2012 and [65] Jasper et al 2012, C4 plants produced carbon dioxide fixation is a 4-carbon compound,

C4 plants have a uniform mesophyll layer with developed bundle sheath around each vein, water use efficiency of the plants [66] Stephen 2014. *In vitro* plants are photomixotrophic, low rates of photosynthesis with low light intensities, low CO₂ and high amount of sucrose in the medium which inhibits photosynthesis [67] Reuther 1991), Rubisco and PEPC were greater in the plantlets have 3-4 leaves than *in vitro* date palm [68] Masmoudi et al. 1999, increasing of FBPase and Rubisco when transition from photoheterotrophic to photo-autotrophic conditions of Banana plants [69] Regev et al. 1997, photosynthesis rates improve when cultures are grown under elevated CO₂ levels [70] Mitra et al. 1998, *in vitro* conditions led to limit leaves photosynthetic incorporation of CO₂, after transplanting tobacco plants to natural conditions reached to normal values [71] Katya et al. 1996, lack of photosynthetic capacity is the main factors responsible for decreasing survival and slow growth of microcuttings [72] Carmen et al. 2007, [73] Debergh et al. 2000, *in vitro* condition the lowest intensities of light, little concentrations of CO₂ and inhibition of photosynthesis led to lower net photosynthetic and dark respiration rate than *ex vitro* plants [74] Slavtcheva and Dimitrova 2001 and [75] Zhang et al. 2009, *in vitro* plantlets detect degradation of Rubisco. In *ex vitro* plantlets, these degradation were no longer detected, decrease in soluble sugars and starch gave an indication of a faster acquisition of autotrophic characteristics [76] Carvalho et al. 2005, at the beginning of acclimatization low vitality *in vitro* plants were noted and indicate poor functioning of dark phase of photosynthesis, vitality of new leaves reflected to higher photosynthetic potential [77] Fuentes et al., 2005, [78] LLisandra et al. 2006 on *Malus domestica* – cv M9 and [79] Pedroso et al. 2010.

V. CONCLUSION

Date palm *in vitro* tissues showed decreasing leaves contents of chlorophyll which as strong indicator of photosynthesis process and its components as proteins and indoles in addition to many enzymes as antioxidant enzymes and the main photosynthesis enzymes for date palm as PEPC and Rubisco that reflected on the growing plants, these effects can be modified after acclimatization in the greenhouse, as well as the plantlets derived tissue culture technique exhibited normal process in the open field.

REFERENCES

- [1] Darwesh, Rasmia, S.S.(2015): Morphology, physiology and anatomy *in vitro* affected acclimatization *ex vitro* date palm plantlets: A Review. International Journal of Chemical, Environmental & Biological Sciences (IJCEBS) 3, 2: 175-182
- [2] Pospisilova', J. C_atsky', Z. S_esta'k, (1997): Photosynthesis in plants cultivated *in vitro*, in: M. Pessaraki (Ed.), Handbook of Photosynthesis, Marcel Dekker, New York, , pp. 525–540.
- [3] Kozai, T., Kubota, C., Jeong, B.R., (1997): Environmental control for large-scale production of plants through *in vitro* techniques. Plant Cell Tissue Org. Cult. 51, 49–56.
- [4] DelaViña G, Pliego-Alfaro F, Driscoll SP, Mitchell V, Parry MA, Lawlor DW (1999): Effects of CO₂ and sugars on photosynthesis and composition of avocado leaves grown *in vitro*. Plant Physiol Biochem 37: 1–9.
- [5] Desjardins Y (1995): Photosynthesis *in vitro* – on the factors regulating CO₂ assimilation in micropropagation systems. Acta Hort 393: 45– 61
- [6] Kozai T, Smith MAL (1995): Environmental control in plant tissue culture general introduction and overview. In: Goto E, Kurata K, Hayashi M, Saa S (eds) Plant Production in Closed Ecosystems. Kluwer Academic Publishers, Dordrecht–Boston–London, pp 153–169
- [7] Rival AK, Triques K, Beule T, Nato A, Lavergne D, Santamaria J, Verdeil JL, Hoche V, Oropeza C, Hamon S (1999): A multi-parameter approach for the study of *in vitro* photosynthesis. Current Plant Sci Biotech Agric 36: 437–440
- [8] Premkumar, A., José A. Mercado, Miguel A. Quesada (2001): Effects of *in vitro* tissue culture conditions and acclimatization on the contents of Rubisco, leaf soluble proteins, photosynthetic pigments, and C/N ratio. Journal of Plant Physiol. 158. 835–840.
- [9] Cournac L, Cirier I, Chagvardieff P (1992): Improvement of photoautotrophic *Solanum tuberosum* plantlet culture by light and CO₂: differential development of photosynthetic characteristics and varietal constraints. Acta Hort 319:53-58
- [10] Kadlecěk, P, Ingrid Ticha' a, Daniel Haisel b, Ve'ra C_ apkova' b, Christian Scha'fer (2001): Importance of *in vitro* pretreatment for *ex vitro* acclimatization and growth. Plant Science 161: 695–701
- [11] Van Huylenbroeck JM, Debergh PC (1996): Physiological aspects in acclimatization of micropropagated plantlets. Plant Tissue Culture and Biotech 2: 137–141.
- [12] Ticha, I; F. C_ a'p, D. Pacovska', D. Haisel, V. C_ apkova', C. Scha'fer, (1998): Culture on sugar medium enhances photosynthetic capacity and high light resistance of plantlets grown *in vitro*, Physiol. Plant. 102 155–162.
- [13] Amâncio S, Rebordão JP, Chaves MM (1999): Improvement of acclimatization of micropropagated grapevine: Photosynthetic competence and carbon allocation. Plant Cell Tiss Org Cult 58: 31–37
- [14] Taiz, L., Zeiger, E. (2010): Plant Physiology. 5th Ed. Sinauer Associates, Sunderland .
- [15] Agetsuma M1, Furumoto T, Yanagisawa S, Izui K. (2005): The ubiquitin-proteasome pathway is involved in rapid degradation of phosphoenolpyruvate carboxylase kinase for C4 photosynthesis. Plant Cell Physiol. 46 (3): 389-398
- [16] Méndez-Lucas A., Hyroššová P., Novellasdemunt L., Viñals F. and Perales J.C. (2014): "Mitochondrial phosphoenolpyruvate carboxykinase (PEPCK-M) is a pro-survival, endoplasmic reticulum (ER) stress response gene involved in tumor cell adaptation to nutrient availability". J. Biol. Chem. 289 (32): 22090–102
- [17] Christopher J.T. and Holtum J.A.M. (1996): "Patterns of carbon partitioning in leaves of Crassulacean acid metabolism species during deacidification". Plant Physiol. 112 (1): 393–399.
- [18] Kanai R. and Edwards G.E. (1999): "3. The Biochemistry of C4 Photosynthesis". In Sage RF, Monson RK. C4 Plant Biology. pp. 43–87.
- [19] Kriengkrai MB, Suriyan C and Chalernpol K (2004): Enhanced growth and photosynthesis of rain tree (*Samanea saman* Merr.) plantlets *in vitro* under a CO₂-enriched condition with decreased sucrose concentrations in the medium. Scientia Horticulturae 103: 51–63.
- [20] Abd El-Baky M.A. (2012): Using Morphological and Anatomical Features as Taxonomical Evidences to Differentiate Between Some Soft and Semi-Dry Egyptian Cultivars of Date Palm. Journal of Horticultural Science & Ornamental Plants 4 (2): 195-200.
- [21] Anushi A. J. and Anis, M. (2014): Changes in Antioxidative Enzymatic Responses during Acclimatization of *In Vitro* Raised Plantlets of *Cardiospermum halicacabum* L. against Oxidative Stress. Plant Physiol Pathol, 2:4: 1-5
- [22] Lichtenthaler H.K. and Welburn E.R. (1994): Determination of total carotenoids and chlorophylls a and b of leaf extracts in different solvents. J. Biochem Soc Tran. 603: 591-593.
- [23] Bradford MM (1976): A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. Anal Biochem 72: 248–254.



- [24] Larsen P, Harbo A, Klungroun, S, Ashein T (1962): On the biogenesis of some indole compounds in *Acetobacter xylinum*. *Physiol Plant*, 15 : 552 – 565
- [25] Aebi, H. (1984): Catalase *in vitro*. *Methods in Enzymology* 105, 121-126.
- [26] Polle, A.; Otter, T. and Seifert, F. (1994): Apoplastic peroxidase and lignification in needles of Norway (*Picea abies* L.). *Plant Physiol*. 106, 53-50
- [27] Gonzalez, M.C., Osuna, L., Echevarria, C., Vidal, J., and Cejudo, F.J. (1998): Expression and localization of phosphoenolpyruvate carboxylase in developing and germinating wheat grains. *Plant Physiology* 116, 1249 – 1258
- [28] Lilley, R.Mc. and Walker, D. A. (1974): An improved spectrophotometric assay for ribulose bis-phosphate carboxylase. *Biochim. Biophys. Acta*. 358: 226-229.
- [29] Snedecor, G.W. and Cochran W.G. (1990): *Statistical Methods*. 11th Ed. Iowa State Univ., Press. Ames, Iowa, USA
- [30] Jain SM (2012): Date palm biotechnology: Current status and prospective-an overview. *Emir J Food Agric* 24: 386-399
- [31] FAO. *Statistical Databases* (2013): www.FAO.org Accessed 20.01.
- [32] Eke, C.R. Akomeah P and Asemota O (2005): Somatic embryogenesis in date palm (*Phoenix dactylifera* L.) from apical meristem tissues from 'zebia' and 'loko' landraces. *African Journal of Biotechnology*, 4 (3): 244-246
- [33] Al-Mazroui H, Zaid A, Bouhouche N. (2007): Morphological abnormalities in tissue culture-derived date palm (*Phoenix dactylifera* L.). III International Conference on Date Palm. *Acta Hort*. 736: 329-335.
- [34] Al-Khateeb AA (2008b): The problems facing the use of tissue culture technique in date palm (*Phoenix dactylifera* L.). *Sci J King Faisal Univ* 9: 85-104.
- [35] Mustafa, N.S., Taha, R.A., Hassan, S.A.M., Ziad, N.S.M. and Mustafa, E.A. (2013): Overcoming phenolic accumulation of date palm *in vitro* culture using α -tocopherol and cold pre-treatment. *Middle-East Journal of Scientific Research* 15 (3): 344-350
- [36] Mazri MA and Meziani R (2015): Micropropagation of Date Palm: A Review. *Cell & Developmental Biology*, 4 (3): 1-5
- [37] Fila, G., Ghashghaie, J., Hoarau, J., Cornic, G. (1998): Photosynthesis, leaf conductance and water relations of *in vitro* cultured grapevine rootstock in relation to acclimatization. - *Physiol. Plant*. 102: 411-418.
- [38] Bolar, J.P., Norelli, J.L., Aldwinckle, H.S., Hanke, V. (1998): An efficient method for rooting and acclimation of micropropagated apple cultivars. - *HortScience* 37: 1251-1252.
- [39] Krause GH and Weis E (1991): Chlorophyll fluorescence and photosynthesis: the basics. *Annu. Rev. Plant Physiol. Plant Mol. Biol.* 42: 313-349
- [40] Carvalho L.C., Os'orio M.L., Chaves M.M. and Am'ancio S. (2001): Chlorophyll fluorescence as an indicator of photosynthetic functioning of *in vitro* grapevine and chestnut plantlets under *ex vitro* acclimatization. *Plant Cell, Tissue and Organ Culture* 67: 271-280
- [41] Bořena Matysiak (2004): Effect of light intensity on growth and chlorophyll fluorescence of *Rhododendron* microcuttings during acclimatization. *Folia Horticulturae Ann.* 16(1): 107-114
- [42] Arigita L, Fernández B, González A, Tamés RS (2005): Effect of the application of benzyladenine pulse on organogenesis, acclimatization and endogenous phytohormone content in kiwi explants cultured under autotrophic conditions. *Plant Physiol Biochem* 43: 161-167.
- [43] Lichtenthaler HK1, Ac A, Marek MV, Kalina J, Urban O. (2007): Differences in pigment composition, photosynthetic rates and chlorophyll fluorescence images of sun and shade leaves of four tree species. *Plant Physiol Biochem*. Aug;45(8):577-88
- [44] Lichtenthaler HK1, Babani F, Navrátil M, Buschmann C. (2013): Chlorophyll fluorescence kinetics, photosynthetic activity, and pigment composition of blue-shade and half-shade leaves as compared to sun and shade leaves of different trees. *Photosynth Res.*, 117(1-3):355-366
- [45] Suthar, R.K. and Purohit S.D. (2011): Biopriming of micropropagated *Boswellia serrate* Roxb. Plantlets – Role of endohytic root fungus *Piriformospora indica*. *Indian Journal of Biotechnology*, 11 (7): 304-308
- [46] Dragolova D., Stefanova M., Dimitrova M., Koleva D., Zhiponova M. and Kapchina – Toteva V. (2015): *In vitro* cultivation and *ex vitro* adaptation of *Nepeta nuda* ssp. *nuda*-correlation between regeneration potential, leaf anatomy and plastid pigments. *Bulgarian Journal of Agricultural Science*, 21 (5): 1027 – 1032.
- [47] Akram M and Aftab F (2015): Effect of Cytokinins on *In vitro* seed Germination and Changes in Chlorophyll and Soluble Protein Contents of Teak (*Tectona grandis* L.). *Biochemistry & Physiology*, 4(3): web page
- [48] Hofman P, Haisel D, Komenda J, Vagner M, Ticha I, Scafer C and Capkova V. (2002): Impact of *in vitro* cultivation on stress responses and on changes in thylakoid membrane proteins and pigments of Tobacco during *ex vitro* acclimation. *Biologia Plantarum*, 45 (2): 189-195.
- [49] Sharifi G (2012): Plant Antioxidative Enzymes –Case Study: *In Vitro* Organogenesis of Saffron (*Crocus sativus* L.). *Department of Basic Sciences, Iranian Encyclopedia Compiling Foundation, Tehran, Iran, INTECH*, 369-380
- [50] Darwesh Rasmia S.S., Zeinab, E. Zaid and Rehab, A. Sidky (2011): Effect of Ammonium Nitrate and GA3 on Growth and Development of Date Palm Plantlets *in Vitro* and Acclimatization Stage. *Research Journal of Agriculture and Biological Sciences*, 7(1): 17-22
- [51] Ibrahim, A.E.; Mona, H.M. and Darwesh, R.S.S (2012): Pre-acclimatization stage of date palm (*Phoenix dactylifera*) plantlets cv. Barhee as affected by potassium nitrate and sucrose. *Journal of Biological Chemistry & Environmental Science*, 7 (3):1-20
- [52] Young I. S. and Woodside J. V. (2001): Antioxidants in health and disease. *J. Clin. Pathol.* 54:176-186.
- [53] Karou D., Dicko M. H., Simpore J. & Traore A. S. (2005): Antioxidant and antibacterial activities of polyphenols from ethnomedicinal plants of Burkina Faso. *Afr. J. Biotechnol.* 4:823-828
- [54] Van Huylenbroeck JM, Piqueras A, Debergh PC (2000): The evolution of photosynthetic capacity and the antioxidant enzymatic system during acclimatization of micropropagated *Calathea* plants. *Plant Sci* 155: 59-66.
- [55] Faisal M, Anis M (2009): Changes in photosynthetic activity, pigment composition, electrolyte leakage, lipid peroxidation and antioxidant enzymes during *ex vitro* establishment of micropropagated *Rauwolfia tetraphylla* plantlets. *Plant Cell Tiss Org Cult* 99: 125-132
- [56] Faisal M, Anis M (2015): Effect of light irradiations on photosynthetic machinery and antioxidant enzymes during *ex vitro* acclimatization of *Tylophora indica* plantlets. *J Plant Interact* 5: 21-27.
- [57] Starlin T. & Gopalakrishnan V. K. (2013): Enzymatic and non-enzymatic antioxidant properties of *Tylophora pauciflora* Wight and Arn – An *in vitro* study. *Asian J. Pharm. Clin. Res.* 6:152-157.
- [58] Agnieszka W and Edyta S (2014): Effect of cytokinins on antioxidant enzymes in *in vitro* grown shoots of *Pelargonium hortorum* L. H. Bayley. *Acta Agrobotanica*, 67 (4): 33-42
- [59] Konieczny R, Banaś A.K., Surówka E, Michalec Z., Miszański Z., and Konieczny M.L. (2014): Pattern of antioxidant enzyme activities and hydrogen peroxide content during developmental stages of rhizogenesis from hypocotyl explants of *Mesembryanthemum crystallinum* L. *Plant Cell Rep.* 2014; 33(1): 165-177.
- [60] Oberhuber W, Dai Z-Y, Edwards GE (1993): Light dependence of quantum yields of Photosystem II and CO2 fixation in C3 and C4 plant. *Photosynthesis research*. 35: 265-274
- [61] Wang C, Longyun Guo, Yixue Li Email author and Zhuo Wang (2012): Systematic Comparison of C3 and C4 Plants Based on Metabolic Network Analysis. *Proceedings of the 23rd International Conference on G*
- [62] Kwa SH, Wee YC and Kumar PP (1997): Ribulose-1, 5-bisphosphate carboxylase and phosphoenolpyruvate carboxylase activities of photoautotrophic callus of *Platyserim coronariuin* (Koenig ex OF Muell.) Desv. under CO2 enrichment. *Plant Cell Tiss. Org. Cult.* 50: 75-82 *enome Informatics (GIW 2012)*
- [63] Doubnerová, V., Ryšlavá, H. (2011) : What can enzymes of C4 photosynthesis do for C3 plants under stress? – *Plant Sci.* 180: 575-583.

- [64] Freschi, L., Mercier, H. (2012): Connecting environmental stimuli and crassulacean acid metabolism expression: Phytohormones and other signaling molecules. – Prog. Bot. 73: 231-255.
- [65] Jasper J.L. Pengelly, Jackie Tan, Robert T. Furbank and Susanne von Caemmerer (2012): Antisense Reduction of NADP-Malic Enzyme in *Flaveria bidentis* Reduces Flow of CO₂ through the C₄ Cycle. *Plant Physiology*, 160 (2): 1070-1080
- [66] Stephen G. S. (2014): Autotrophism: Carbon Reactions (Calvin Cycle, C₄, CAM). *Plant Physiology (Biology 327)*: web page
- [67] Reuther G (1991): Stimulation of photoautotrophy of in vitro plants. *Acta Hort.* 300: 59-75.
- [68] Masmoudi R., Rival A, Nato A, Lavergne D, Drira N and Mazri G, M.A. and Meziani R (2015): Micropropagation of Date Palm: A Review. *Cell & Developmental Biology*, 4(3): 1-5
- [69] Regev, I.; Gepstein, S.; Duvdevani, A.; Magdar, D.; Khayat, E.; Meristem, R.; Hanikra, R. (1997): Carbon Assimilation, Activities of Carboxylating Enzymes and Changes in Transcript Level of Genes Encoding Enzymes of the Calvin Cycle and Carbon Partitioning of Banana Plantlets during Transition from Heterotrophic to Autotrophic Existence. *Acta horticulturae*, 447: 561-568.
- [70] Mitra A, Bhattacharya PS, Dey S, Sawarkar SK, Bhattacharyya BC (1998): Photoautotrophic in vitro culture of *Chrysanthemum* under CO₂ enrichment. *Biotechnol Techn* 12:335-337
- [71] Katya G., Ivan Y., and Antoaneta K. (1996): Photosynthetic characteristics of transformed Tobacco plants grown in vitro after their transplantation in natural conditions. *Bulg. Journal Plant Phtsiol.*, 22 (3-4): 3-13
- [72] Carmen V-A, Sandra B. W., Michael E. K., and Nancy L. P. (2007): Influence of in vitro growth conditions on in vitro and ex vitro photosynthetic rates of easy- and difficult-to-acclimatize sea oats (*Uniola paniculata* L.) genotypes. *In Vitro Cellular and Developmental Biology - Plant* 43(3):237-246.
- [73] Debergh P.C., Topoonyanont N., Vanhuylenbroeck J.M., Moreir H., Oysert E., (2000): Preparation of microplants for ex vitro establishment. *Acta Hort.* 530: 269-273
- [74] Slavtcheva, T. and Dimitrova, V. (2001): Gas exchange of in vitro and ex vitro grown grapevine plants. *Photosynthetica*, Prague, 39: 29-33
- [75] Zhang M., Zhao D., Ma Z. and Xiao X.Y. (2009): Growth and Photosynthetic Capability of *Momordica grosvenori* Plantlets Grown Photoautotrophically in Response to Light Intensity. *HortScience*, 44 (3): 757-763
- [76] Carvalho L.C., Esquível M.G. and Amâncio S. (2005): Stability and Activity of Rubisco in Chestnut Plantlets Transferred to Ex vitro Conditions under Elevated CO₂. *In Vitro Cellular & Developmental Biology. Plant*, 41 (4): 525-531
- [77] Fuentes G, Talavera C, Desjardins Y, Santamaria JM (2005): High irradiance can minimize the negative effect of exogenous sucrose on the photosynthetic capacity of in vitro grown coconut plantlets. *Biol. Plant.* 49: 7-15.
- [78] Iliandra Z; Marcos A B; Douglas D S; Eugenia J B B; José A P (2006): Chlorophyll fluorescence in in vitro cultivation *Apple. R. Bras. Agrobiência, Pelotas*, v. 12, n. 3, p. 305-308
- [79] Pedroso A.N.V.; Lazarini R.A.M.; Tamaki V. and Nievola C.C. (2010): In vitro culture at low temperature and ex vitro acclimatization of *Vriesea inflata* an ornamental bromeliad. *Brazilian Journal of Botany*, 33 (3): web page




Fig. 1. Different type's tissues affected chlorophyll contents




Fig. 2. Different type's tissues affected proteins contents




Fig. 3. Different type's tissues affected proteins contents




Fig. 4. Different type's tissues affected CAT and POD activity




Fig. 5. Different type's tissues affected PEPC activity




Fig. 6. Different type's tissues affected Rubisco activity



Fig. 7. In vitro and ex vitro date palm stages. In vitro stages as 1= shooting stage 2 = rooting stage). Ex vitro stages as 3 and 4 = plantlets in the greenhouse 5 = plantlets in the open field 6 = offshoot.