

Does Climate Change Allow Grapevine Growing in the Southernmost Finland

Juha Karvonen

Department of Agricultural Sciences,
 Faculty of Agriculture and Forestry, University of Helsinki, Finland
 Email: juha.i.karvonen@helsinki.fi

Abstract – According to the scenarios estimating climate change, the annual mean air temperatures could rise from the current by 3°C to 4°C in the Helsinki area, and the growing season will become 30 to 40 days longer by the year 2100. The purpose of this study was to establish how much the annual mean air temperature between the years 2005 to 2014 have risen in the Helsinki area and how much the growing season has been extended in comparison to the mean in the years from 1971 to 2000. The results indicate that in the Helsinki region, the annual mean air temperature has risen by 1.1°C, and growing season has extended by 27 days between the years 2005 and 2014, as compared to the 1971 to 2000 levels. Over the past ten-year period, the climatic temperature and the length of the growing season during the summer months in the southernmost Finland have moved closer to the climate and growing season of Central Europe, and thanks to this, it is possible to grow a harvestable grapevine in the southernmost regions of Finland.

Keywords – Air Temperature, Climate Change, Growing Degree Days, Growing Season, Nordic Grapevine Growing.

I. INTRODUCTION

The increasing levels of greenhouse gasses have caused the raise of earth's surface temperatures in the world. During the past 100 years, the annual mean air temperature in the southernmost parts of Finland (60°N; 24°E) has risen by 1°C [1]. The current ever-increasing usage of fossil fuels has increased the global carbon dioxide concentration which has accelerated global warming together with the increase in the water vapour concentration in the air caused by this warming [2]. According to the Rosby Centre's regional climate model, these greenhouse gases are predicted to raise the mean yearly temperatures of the Northern European climate by +4°C by the year 2100. This increases in temperatures at its greatest during the winter months in particular [3]. The scenarios of the IPCC (Intergovernmental Panel on Climate Change) for the years from 1996 to 2013 predict

similar changes, if carbon-dioxide emissions are not greatly reduced.

In Finland, a warming of the climate has been detectable since the years 1971 - 2000 as an increase in the annual mean air temperatures and an extension of growing season [4]. At the same time, there has been a significant increase in temperature sums of growing degree days +5°C and +10°C [5]. This has introduced to the southern regions of Finland and the other Nordic countries some new cultivated plant species for experimentation, such as corn and grapevine. In these regions, interest towards wine growing has increased year by year. The total wine growing area in the Nordic Countries is estimated to be 300 to 500 hectares. Sweden and Denmark are already listed as EU wine-producing countries, and in Finland will also become a wine-producing country in the coming years.

The purpose of this study was to examine how much the climate and grapevine growing conditions, start of growth, and harvest ripening in Helsinki region differ from the growing conditions and growth cycle in Central Europe at the present time.

II. MATERIAL AND METHODS

The study was undertaken in three locations in Central and in one location in Northern Europe (Table 1). It studied the changes in the climatic temperature and growing season length in the Helsinki region over the past 15 years. In addition, a comparison was made of the air and soil temperatures in the Helsinki region from May to October with the temperatures of the three Central European areas, and the development of grapevine sprouts and blossoms was observed in these areas based on phenological observations (12 leaves separated, inflorescence well developed, single flowers separated; E-L number 17) [6], and harvest maturing was observed by measuring total grape sugar concentrations.

Table 1: Observation locations and grape varieties

| Community | Country | Location | Above sea level (m) | Distance to Herrlisheim-prés-Colmar (km) | Grape variety |
|-------------------------|---------|------------------|---------------------|--|---------------|
| Herrlisheim-prés-Colmar | France | 48°01'N, 07°19'E | 242 | 0 | Pinot Noir |
| Freyburg, Unstrut | Germany | 51°13'N, 11°46'E | 110 | 360 | Regent |
| Neubrandenburg | Germany | 53°33'N, 13°16'E | 53 | 550 | Regent |
| Helsinki region | Finland | 60°24'N, 25°01'E | 63 | 1390 | Rondo |

Observations on air temperatures were based on the statistical data of the Finnish Meteorological Institute and Meteorological ServisForeca (www.foreca.com). Soil

temperature measurements were taken at depths of 20 cm and 40 cm using digital thermometers that recorded maximum and minimum temperatures (Waterproof In-Out

Door Max-Min Thermometer with Hygrometer, Shenzhen Hong Tong Yuan Technology Ltd, China, Shenzhen). The total sugar content of the grapes ($^{\circ}\text{Bx}$) was measured using a Brix refractometer (HR-180, Optika, Ponteranica, Italy). A statistical treatment was completed with a t-test: $t = \frac{\bar{X}-\bar{Y}}{\sqrt{V}}$.

As part of the t-test, \bar{X} was the mean value of the variables, and \bar{Y} was the mean value of the other variables, and V an estimate of the variance of random error. $P < 0.05$ was used to estimate statistical significance

III. RESULTS AND DISCUSSION

The average temperature of the globe varies yearly approximately by a tenth of a degree. Disregarding this

variation, the average temperature of the Earth has risen since 1880s approximately by 0.85°C [7]. This has occurred in the northern parts of the Europe as well. The temperature of the Finnish climate over an observation period of 160 years has increased in the 21st century by 0.6°C in comparison to the mean of 160 years, and this increase has been most obvious during spring [7]. Tables 2 and 3 show that the yearly averages of the temperatures of the ten-year periods between 1971 to 2010 have constantly risen, and that the temperature averages of single years during the last ten-year period (2005 to 2014) have risen by 1.1°C in comparison to the average of years 1971 to 2000; this supports the results published by Tuomenvirta [8].

Table 2: Mean values of air temperatures in the Helsinki region over a 40-year period (1971 – 2010) and the ten-year period (2005 – 2014)

| Periods of ten years 1971 – 2010 | Mean air temperatures ($^{\circ}\text{C}$) | Years 2005 – 2014 | Mean air temperatures ($^{\circ}\text{C}$) |
|-------------------------------------|--|-------------------|--|
| 1971 – 1980 | 5.0 | 2005 | 6.6 |
| 1981 – 1990 | 5.3 | 2006 | 6.7 |
| 1991 – 2000 | 5.4 | 2007 | 7.0 |
| 2001 – 2010 | 6.0 | 2008 | 7.6 |
| 2011 – 2020 ¹⁾ | 6.5 | 2009 | 6.2 |
| 2021 – 2030 ¹⁾ | 6.8 | 2010 | 5.1 |
| 2031 – 2040 ¹⁾ | 7.2 | 2011 | 7.2 |
| 2041 – 2050 ¹⁾ | 7.5 | 2012 | 6.2 |
| 2070 – 2080 ¹⁾ | 8.0 | 2013 | 6.6 |
| 2090 – 2100 ¹⁾ | 8.8 | 2014 | 7.4 |

Finland is the northernmost agricultural country in the world, and due to its northern location, average temperatures in Finland vary more from one year and one decade to the next, than the global average temperatures. One reason for this are the low pressure variations coming from the North Atlantic. The low pressure areas that come to Finland via the northern route will raise the yearly temperature, but the low pressures arriving to the southern parts are replaced by easterly winds bringing cold air, which cause the temperature to drop [7]. The noticeable increase by 1.1°C in the Table 3 was apparently caused by a large deviation, and was not statistically significant ($P > 0.05$).

The temperature of the Finnish climate is predicted to rise by 2°C by the year 2050, and by 3°C by the year 2080, at the same time as rainfall increases up to 30% [8]. Such a significant warming of the climate will allow for

the profitable cultivation of plants thriving in mild climates, such as the grapevine, in southern Finland. Even today, a few hybrid vines will have time to ripen well in the Helsinki area [10]. Should the yearly average temperature of southern Finnish climate rise 2°C by the year 2050, the yearly mean temperature of the Helsinki region will reach 8°C to 9°C in 40 years, *i.e.*, close to the temperatures of the current wine growing regions of Central Europe. Then it will be possible to grow regular fast ripening varieties, such as Chardonnay and Pinot noir [11]. In the new growth condition, - thanks to the long day, humus-rich soil, and sufficient rainfall - the grapevine could produce even more abundant and superior quality crops than in the current wine-growing areas, where heat and drought would, according to forecasts, increase and hamper farming.

Table 3: Changes of air temperatures and lengths of growing seasons in the Helsinki region during the ten-year period (2005 – 2014) compared to the 30-year period (1971 – 2000).

| | | | |
|---|---|--|---|
| Mean air temperature 1971 – 2000; 5.6 ± 0.4 ($^{\circ}\text{C}$) | Mean air temperature 2005 – 2014; 6.7 ± 0.7 ($^{\circ}\text{C}$) | Difference of mean values; 1.1 ($^{\circ}\text{C}$) | Statistical significance $t=1.5607$; $P > 0.05$ |
| Mean length of growing season 2005 - 2014 198 ± 18 days | Mean length of growing season 1971 - 2000 171 ± 15 days | Difference of mean values 27 days | Statistical significance $t=5.5214$; $P < 0.05$ |

In this study, the lengthening of the growing season in the Helsinki region was demonstrated more clearly than the increase in climatic temperature. In comparison to the

lengths of the growing seasons between 1971 and 2000, the growing periods' mean values during the past ten-year period have lengthened by 27 days, which is statistically

significant ($P < 0.05$). The lengthening of the growing period has manifested itself in earlier springs [7]. According to Bauer [12], the grapevine requires, depending on variety, a growing season of approximately 200 days. Table 3 shows that the mean growing season

length in the Helsinki region has been 198 days in the years 2005 to 2014, in 6 of these years it has been over 200 days [5], which should suffice for cultivating some Central-European *Vitisvinifera* varieties.

Table 4: Mean monthly air temperatures

| Locality | May | June | July | August | September | October |
|----------------|----------|----------|----------|----------|-----------|----------|
| Helsinki | 10.9±4.2 | 13.9±4.6 | 19.8±3.0 | 17.2±3.9 | 12.1±3.3 | 5.7±4.3 |
| Herrlisheim | 14.3±2.9 | 18.6±3.2 | 20.2±2.5 | 17.7±2.7 | 15.1±2.4 | 12.5±2.9 |
| Freyburg | 12.5±3.6 | 16.6±4.1 | 20.0±2.8 | 16.9±2.9 | 15.8±2.6 | 12.6±2.8 |
| Neubrandenburg | 12.1±3.4 | 15.7±3.1 | 20.0±2.6 | 16.7±2.9 | 15.1±2.4 | 11.6±2.6 |

Tables 4 and 5 compare the six-month temperatures of the Helsinki region growing season to the temperatures of the three Central European comparison regions. It shows that the mean monthly temperatures in July and August in the Helsinki region are statistically not significantly different from those of the comparison regions ($P > 0.05$). During other months, the differences between the temperatures of the Helsinki region and other regions are

statistically significant, with the exception of northern Neubrandenburg, where wine growing is being restarted. In 2014, June was exceptionally cold in southern Finland, but July and August were sweltering, as usual. During July and August, there were 44 days of hot weather, when the highest temperature of the day exceeded 25°C and of which in 22 days temperatures exceeded 35°C [13].

Table 5: Statistical significance of mean air temperatures between localities during three midsummer months

| Month | n | Helsinki >>Herrlisheim | Helsinki >>Freyburg | Helsinki ><Neubrandenburg |
|--------|----|------------------------|---------------------|---------------------------|
| June | 30 | t=4.5988; P<0.05 | t=2.4107; P<0.05 | t=1.7822; P>0.05 |
| July | 31 | t=0.5714; P>0.05 | t=0.2702; P>0.05 | t=0.2857; P>0.05 |
| August | 31 | t=0.6849; P>0.05 | t=0.3448; P>0.05 | t=0.6579; P>0.05 |

n = number of days

In late May, early June of 2014 (25 May to 3 June), soil temperatures measured in Herrlisheim-prés-Colmar at depths of 20 cm and 40 cm were at least 2.5°C higher than in other regions, due to the earlier spring and shorter duration or lacking of snow coverage. Moving northward, the temperatures of soil were lower due to the later than

usual spring and warming of air temperature. In Helsinki region, the soil temperatures were clearly lower, even on 3 June than in other regions. Compared with the soil temperatures in Neubrandenburg, the differences at 20 cm and 40 cm depths were clearly the lowest as compared with others, i.e., 0.6°C to 1.2°C (Table 6).

Table 6: Mean soil temperatures at the depths of 20 cm and 40 cm, grapevine growing stage (Eichhorn-Lorenz), total sugar content of grapes (°Bx) and measurement dates

| Locality | Depth of 20 (cm); (°C) | Depth of 40 (cm); (°C) | Dates for temperatures | Eichhorn-Lorenz: 17 | (°Bx) | Dates for (°Bx) |
|----------------|------------------------|------------------------|------------------------|---------------------|----------|-----------------|
| Helsinki | 13.6 | 12.0 | 3 June | 4 June | 16.6±0.5 | 2 October |
| Herrlisheim | 17.0 | 16.6 | 25 May | 26 May | 17.9±1.0 | 7 September |
| Freyburg | 14.5 | 13.8 | 30 May | 1 June | 13.1±0.8 | 9 September |
| Neubrandenburg | 14.2 | 13.2 | 1 June | 1 June | 13.0±1.3 | 9 September |

Grapevine growth stage 17 [6] (Eichhorn and Lorenz 1977), which is equivalent to the BBCH scale 55 [14], was reached in the Helsinki region 9 days later than in Herrlisheim, and just 3 days later than in Freyburg and Neubrandenburg. Although during the vines' sprouting and inflorescence stages producing red wine grapes the regional differences were only 3 to 7 days, at the harvesting stage they were several weeks, based on the grape total sugar level (°Bx); this may be due at least in part to both climatic differences and different grape varieties (Tables 1 and 6). Based on observations, only a 3 to 4 days late blooming in the spring may delay even during a normal summer the ripening and harvesting of the grapes by 1 to 2 weeks in the autumn. During colder than

usual summers, ripening of grapes may be delayed to even later date.

During the last 40 years, the temperatures of the Helsinki region have risen over 1°C which is the equivalent of a lengthening of the growing season by 10 to 11 days [2]. The temperature increase has been most obvious during the past 10 years. By the end of the century, and with the realisation of the IPCC A1B scenario, thermal growing seasons will get longer during the autumn and spring in the whole of Finland by a total of 35 to 40 days [15]. When at the same time snow coverage will diminish, the climates of southern parts of Finland and the other Nordic countries will change to resemble the current climate in central Europe [16].

IV. CONCLUSIONS

In Finland, a significant warming of the climate has been observed during the last decades. This has encouraged to try out some new cultivated species as grapevine, in the southernmost Finland in the Helsinki region. Also the results of this study show that in the Helsinki region (and Baltic Sea coast) the annual mean air temperature has risen and the growing season has continually extended. The mean air temperature for two summer months (July and August) did not differ significantly ($p > 0.05$), but the soil temperatures were lower when compared to traditional winegrowing areas of central Europe. Deduced from the measured sugar content of grapes ($^{\circ}\text{Bx}$), the grapes were harvest ripe in the Helsinki region several weeks later than in the other localities. The climate differences between northern and southern localities will probably decrease according to numerous climate scenarios. The existing climate in the southernmost parts of Finland does not prevent grapevines from survival from climate-induced stress, nor viticulture from using suitable varieties, and if the climate continues to warm up, as predicted, the grapevine may become a cultivated plant in the southern Finland already by the year 2040.

AUTHOR'S PROFILE



Dr. Juha Karvonen

has completed his M.Sc. (hortic.) studies from Faculty of Agriculture and Forestry, University of Helsinki in 2009 and started doctoral studies in 2013. Juha Karvonen has published a book on Nordic viticulture in 2004.

REFERENCES

- [1] H. Tietäväinen, H. Tuomenvirta, and A. Venäläinen, "Annual and seasonal mean temperatures in Finland during the last 160 years based on gridded temperature data," *International Journal of Climatology*, vol. 30(15), 2010, pp. 2247-2256.
- [2] IPCC, "The Physical Science Basis. Summary for Policymakers," *Working group I*, 2007.
- [3] P. Samuelsson, C. G. Jones, U. Willen, A. Ullerstig, S. Gollvik, U. Hansson, C. Jansson, E. Kjellström, G. Nikulin, and K. Wyser, "The Rossby Centre Regional Climate model RCA3: model description and performance," *Tellus 63A*, vol. 63, 2011, pp. 1-3.
- [4] Finnish Meteorological Institute, *Newsletter Archive*, 2010.
- [5] J. Karvonen, "The annual growth cycle of grapevines in Southern Finland," *Vitis*, vol. 53(4), 2014, pp.175-180.
- [6] K. W. Eichhorn, and D. H. Lorenz, "Phänologis Entwicklung der Rebe," *Nachrichtenblatten des Deutschen Pflanzenschutzdienstes*, vol. 21, 1977, pp. 119-120.
- [7] T. D. Stoker, D. Qin, G.-K. Plattner, M. Tigor, S. Allen, J. Boschung, A. Nauels, Y. Xia, V. Bex, and P. Midgley, "Climate Change 2013: The Physical Science Basis," Cambridge University Press, 2013, p. 1535.
- [8] H. Tuomenvirta, "Reliable estimation of climatic variation in Finland," *PhD thesis*, University of Helsinki, Helsinki, Finland, 2004.
- [9] IPCC, "Impacts, Adaptation and vulnerability," *Working group II*, 2010.
- [10] J. Karvonen, "Vitis cv. Zilga is a vine for the northern temperate – climate, Short communication," *Horticultural Science (Prague)*, vol. 41(3), 2014, pp.147-151.
- [11] J. Robinson, *The Oxford Companion to Wine*, 3rd edition, Oxford University Press, 2006, pp. 317-324.
- [12] K. Bauer, "Standortansprüche", in *Weinbau*, 7th edition, ÖsterreichischerAgrarverlag, 2002, pp. 124-127.
- [13] Finnish Meteorological Institute, *Statistics*, 2014.
- [14] U. Maier, "Growth stages of mono and dicotyledonous plants", in *BBCH Monograph*, Federal Biological Research Centre for Agriculture and Forestry, Berlin, Germany, 2001.
- [15] L. Neitiniemi-Urpola, "Climate change, threat or opportunity", in *Climate Seminar*, Kajaani, Finland, Finnish Meteorological Institute, 2010.
- [16] IPCC, "Climate change," *Summary for policymakers*, 2014.